

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Agronomy & Horticulture -- Faculty Publications

Agronomy and Horticulture Department

1992

Pathogenicity of *Pseudomonas gladioli* pv. *gladioli* on Rhizomatous Iris and Its Possible Role in Iris Scorch

Annie L. Wrobel

University of Nebraska--Lincoln, awrobel1@unl.edu

John E. Watkins

University of Nebraska-Lincoln, jwatkins1@unl.edu

Don Steinegger

University of Nebraska--Lincoln, dsteinegger1@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/agronomyfacpub>



Part of the [Plant Sciences Commons](#)

Wrobel, Annie L.; Watkins, John E.; and Steinegger, Don, "Pathogenicity of *Pseudomonas gladioli* pv. *gladioli* on Rhizomatous Iris and Its Possible Role in Iris Scorch" (1992). *Agronomy & Horticulture -- Faculty Publications*. 673.

<https://digitalcommons.unl.edu/agronomyfacpub/673>

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Pathogenicity of *Pseudomonas gladioli* pv. *gladioli* on Rhizomatous Iris and Its Possible Role in Iris Scorch

Annie L. Wrobel¹, John E. Watkins², and Donald H. Steinegger³

University of Nebraska, Lincoln, NE 68583-0722

Additional index words. *Iris* spp.

Scorch of rhizomatous iris, *Iris* spp., occurs throughout much of the United States (Black, 1984). It is characterized as a rust-colored browning that starts on the tips of the youngest leaves in the center of the fan. Leaf necrosis progresses downward on the leaves and outward on the fan until the entire fan is affected. Roots shrivel and decay. Root cortical tissue completely deteriorates, but the epidermis and stele remain intact. The rhizome appears unaffected (Wadekamper, 1972).

Because of the sporadic occurrence of scorch and the difficulty of artificially reproducing scorch symptoms, little is known about the etiology, epidemiology, or the pathogenicity of organism(s) involved in its development (Black, 1984). Attempts to identify a predominant causal organism or transmit scorch through inoculation with various fungi and bacteria recovered from scorched iris have failed to reproduce the disease (Wadekamper, 1972).

Bald (1971) implicated the bacterium *Pseudomonas gladioli* pv. *gladioli* (Severini) as the probable cause of iris scorch. He obtained evidence of pathogenicity by inoculating wounded iris tissue with cell suspensions of the *Pseudomonas* bacterium. However, symptoms were less severe than

those that occurred in the field under natural disease development. Rainio (1936) reported that *P. marginata*, synonymous with *P. g. pv. gladioli*, entered the iris leaf through wounds, where it caused a localized water-soaked wet rot but not the characteristic scorch symptoms.

Since the pathogenicity of *P. g. pv. gladioli* to rhizomatous iris is not clear, the present study was undertaken to determine if *P. g. pv. gladioli* could be recovered from scorched iris and to characterize the symptoms produced by inoculation of rhizomatous iris with *P. g. pv. gladioli*.

To determine if *P. g. pv.* was present in scorched iris, isolations were made from symptomatic root and leaf tissues. Excised tissue sections were surface sterilized 2 min in 0.5% sodium hypochlorite and rinsed with sterile distilled water. These were placed in a 125-ml Erlenmeyer flask with 10 ml sterile 12.5 mM PO₄ buffer, pH 7.1, containing 10 mM MgSO₄, and incubated overnight on a rotary shaker. Resulting suspensions were serially diluted in PO₄ buffer and plated onto nutrient broth yeast (NBY) agar medium (Vidaver, 1967). Plates were incubated 2 days at 25 to 27°C. We selected 26 bacterial colonies, based on colony morphology, for SDS-polyacrylamide gel electrophoresis (PAGE) comparison of whole cell polypeptide profiles with those of two reference strains of *P. g. pv. gladioli* (New Zealand Plant Disease Division Cultured Collection Strains 3950 and 3951) (Gwen and Jackman, 1982).

To examine pathogenicity of *P. g. pv. gladioli*, roots, rhizomes, crowns, and leaves of separate plants of 'Victoria Falls' were inoculated with suspensions of *P. g. pv. gladioli* Strains 3950 and 3951 at a concentration of 1×10^8 cfu/ml using a sterile 1.0-cm³ tuberculin syringe. Inoculated plants were placed in a humidity chamber, maintained at 95% to 100% relative humidity and 21 to 30°C for 1 week, after which they were moved to a greenhouse bench where symptom development was monitored. After 8 weeks,

P. g. pv. gladioli was reisolated from lesions and identified using SDS PAGE.

The attempts to recover and identify *P. g. pv. gladioli* from leaf and root tissues of scorched iris were unsuccessful. A comparison of the polypeptide band patterns of the 26 bacterial colonies with those of the reference *P. g. pv. gladioli* strains showed no evidence of *P. g. pv. gladioli* in the tissues examined. Furthermore, inoculation of healthy iris with *P. g. pv. gladioli* Strains 3950 and 3951 failed to produce scorching of inoculated plants. Inoculated roots and rhizomes were asymptomatic. Inoculated crowns sometimes showed signs of localized water soaking at the point of inoculation when some leaf tissue remained; but when leaf tissue was totally removed, inoculated crowns remained symptomless. Leaf inoculations produced irregular, translucent, water-soaked lesions that ranged from 2.4 to 5.6 cm long. The two *P. g. pv. gladioli* strains differed in their virulence with Strain 3950 producing significantly longer lesions than Strain 3951. These data were consistent with those reported by Rainio (1936) but did not confirm the findings of Bald (1971).

Under the conditions of this study, *P. g. pv. gladioli* was not identified among bacteria isolated from scorched iris nor were scorch symptoms induced when healthy iris were inoculated with two strains of *P. g. pv. gladioli*. The pathogenicity of *P. g. pv. gladioli*, previously reported to cause scorch (Bald, 1971), in scorch development remains unclear, as does the etiology of this iris disease.

Literature Cited

- Bald, J.G. 1971. Scorch disease of rhizomatous iris. *Calif. Agr.* 25(2):6-7.
- Black, P. 1984. Scorch: Mystery killer at large. *Amer. Iris Soc. Bul.* no. 252:32-34.
- Gwen, R.J. and P.J.H. Jackman. 1982. The similarities between *P. paucimobilis* and allied bacteria derived from analysis of deoxyribonucleic acids and electrophoretic protein patterns. *J. Gen. Microbiol.* 128:2945-2954.
- Rainio, A.J. 1936. Investigations on bacterial diseases of gladioli (*Pseudomonas marginata* McCl., *P. gummi-sudans* McCl., *Bacillus omnivorus* Hall, and *B. variegatus* Rainio nov. spec.) (German summary). *Valt. Maataloustalk.* 84.
- Vidaver, A.K. 1967. Synthetic and complex media for the rapid detection of fluorescence of phytopathogenic pseudomonas: effect of the carbon source. *Applied Microbiol.* 15:1523-1524.
- Wadekamper, J. 1972. Some research work carried out on the condition of scorch in iris. *Bul. Amer. Iris Soc.* no. 204:21-23.

Received for publication 18 Mar. 1991. Accepted for publication 2 Dec. 1991. This paper is published as Journal Series no. 9528 by the Agricultural Research Div., Institute of Agriculture and Natural Resources, Univ. of Nebraska-Lincoln. This project was funded by a grant from the American Iris Society Foundation. We thank Dan Kuczmarski, Debbie Fujimoto, Monte Miles, and Anne Vidaver for their assistance. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

¹Former Graduate Student, Dept. of Plant Pathology.

²Professor, Dept. of Plant Pathology.

³Professor, Dept. of Horticulture.